



MP Biomedicals, LLC.

29525 Fountain Parkway
Solon, Ohio 44139

Telephone: 440/337-1200

Toll Free: 800/854-0530

Fax: 440/337-1180

mailto:biotech@mpbio.com

web: www.mpbio.com

cappel

**PEROXIDASE-CONJUGATED GOAT AFFINITY PURIFIED
ANTIBODY TO RABBIT IGG F(AB')₂ - 2ML**

Catalog #: 55691

Lot #: 06411

Total Protein: 1.1 mg/ml

Titer: 1:16,000

DESCRIPTION:

Product is horseradish peroxidase (HRP)-conjugated goat affinity purified antibody to rabbit IgG F(ab')₂ and buffer salts.

Antibody and highly purified HRP ($R_z > 3.0$) are conjugated under defined conditions to obtain optimally labeled product. Conjugated protein is purified by salt fractionation. The product is dialyzed into 0.02M sodium phosphate, 0.14M sodium chloride, 1% BSA, pH 7.3, adjusted to standard titer, filtered through a 0.22 μ m filter, vialled and lyophilized. No preservative is added.

USES:

This product is suitable for use as a reagent in enzyme immunoassays (EIA), cell and tissue staining (for light microscopy), cell and tissue labeling (for electron microscopy), and blot immunostaining. Use of affinity purified antibodies is recommended to avoid non-specific binding from the inherent antibodies of host animals. (Note: F(ab')₂ fragments are recommended for staining of cells or tissues which contain Fc receptors. See **Related Product Ordering Information** for these products.)

SPECIFICATIONS:

The total protein is measured using the Biuret procedure with bovine albumin as a standard. To ensure that the product titer falls within the required range, antibody titer is standardized by microtiter plate ELISA with rabbit IgG. The product is tested for purity and specificity at final concentration by immunoelectrophoresis. The antibody is goat IgG; no trace of albumin is detected. It shows reactivity to rabbit F(ab')₂ and IgG; cross-reactivity to other species may exist. Because the product is directed against F(ab')₂, some antibodies are expected to bind light chain sites common to all immunoglobulins. Antibody activity to Fc and non-immunoglobulin serum proteins is not present.

RECONSTITUTION:

Reconstitute product with 2.0 ml of deionized or distilled water. Gentle swirling may be used to speed rehydration. Do not add sodium azide. Avoid vigorous shaking of the reconstituted material.

STORAGE:

Store lyophilized product at 2-8°C. For long-term storage, aliquot and freeze reconstituted product at -20°C or lower.

WARNING AND CAUTION:

This product is intended **FOR RESEARCH USE ONLY**, not for use in diagnostic or therapeutic procedures involving humans or animals.


RELATED PRODUCT ORDERING INFORMATION:

Catalog # Description Size

55604 GOAT ANTI-RABBIT IGG F(AB')₂ 2ML
55624 IGG GOAT ANTI-RABBIT IGG F(AB')₂ 5ML
55637 F(AB')₂ GOAT ANTI-RABBIT IGG F(AB')₂ 2ML
55643 AFF.PUR. GOAT ANTI-RABBIT IGG F(AB')₂ 2MG
59299 AKP AFF.PUR. GOAT ANTI-RABBIT IGG F(AB')₂ 1ML
55678 HRP GOAT ANTI-RABBIT IGG F(AB')₂ 2ML
55687 HRP F(AB')₂ GOAT ANTI-RABBIT IGG F(AB')₂ 2ML
55946 PURIFIED RABBIT IGG F(AB')₂ 10MG
36315 DAB 6x1ML
36317 OPD 6x1ML
36318 TMB 6x1ML
36319 ABTS\ 6x1ML
55984 NORMAL GOAT SERUM 2ML

For fluorochrome conjugates, please see our catalog.

\ABTS is a registered trademark of Boehringer Mannheim GmbH.



Approved by: Christina Marotta
Quality Control Director

**TO PLACE AN ORDER IN THE U.S., PLEASE CALL 800-854-0530
OR TO OBTAIN TECHNICAL ASSISTANCE IN THE U.S., PLEASE CALL 800-279-5490
OUTSIDE THE U.S., CONTACT YOUR LOCAL DISTRIBUTOR.**