



**MP Biomedicals, LLC.**

29525 Fountain Parkway  
Solon, Ohio 44139

Telephone: 440/337-1200

Toll Free: 800/854-0530

Fax: 440/337-1180

mailto:biotech@mpbio.com

web: www.mpbio.com

# cappel

**GOAT AFFINITY PURIFIED ANTIBODY TO  
HAMSTER IGG (WHOLE MOLECULE) - 2MG**

**Catalog #: 56984**

**Lot #: 07336**

Total Protein: 1.0 mg/ml  
Active Antibody: 1.0 mg/ml

## **DESCRIPTION:**

Product is goat affinity purified antibody to hamster IgG (whole molecule) and buffer salts.

The affinity purified antibody is prepared from the specific goat IgG fraction by solid phase absorption. The resulting affinity purified antibody is dialyzed into 0.02M sodium phosphate, 0.14M sodium chloride, pH 7.3, adjusted to standard titer, filtered through a 0.22  $\mu$ m filter, and vialled. No preservative has been added.

## **USES:**

This product is suitable for use as a precipitating antibody. It gives sharp precipitin lines in immunoelectrophoresis and immunodiffusion techniques.

This product is suitable for conjugation with enzymes, radiolabels, fluorochromes or other markers, and for attachment to solid supports for use as an immunoabsorbant.

This product is suitable for use as a primary reagent in enzyme immunoassays, fluorescein immunoassays and blot, cell or tissue immunostaining. These affinity purified antibodies are recommended to avoid non-specific binding from the inherent antibodies of host animals.

See **Related Product Ordering Information.**

## **SUGGESTED WORKING DILUTIONS:**

1:50 - 1:200 for Tissue Staining  
1:100 - 1:500 for Immunoblotting  
1:400 - 2,000 for Elisa

**It is recommended that each lab obtain their own optimal working dilution for each assay**

## **SPECIFICATIONS:**

The total protein is measured by absorbance ( $A_{280nm}$ ). Antibody titer is standardized by equivalence-point precipitation. Each affinity purified antibody is tested for purity at 10 mg/ml using immunoelectrophoresis. The product is goat IgG; no trace of albumin is detected. It shows reactivity to hamster IgG; cross-reactivity to other species may exist. Because the antibody is

directed against whole IgG, some antibodies are expected to bind light chain sites common to all immunoglobulins. Antibody activity to non-immunoglobulin serum proteins is not present.

**EXPIRATION DATE:**

9/30/2018

**STORAGE:**

Store unopened product at 2-8°C. Store opened material no longer than 1-2 days at 2-8°C unless preservative is added. For long-term storage, aliquot and store with 40% glycerol (v/v) at -20°C.

**WARNING AND CAUTION:**

This product is intended **FOR RESEARCH USE ONLY**, not for use in diagnostic or therapeutic procedures involving humans or animals.

**RELATED PRODUCT ORDERING INFORMATION:**

Catalog #   Description   Size

55394	GOAT ANTI-HAMSTER IGG (WHOLE MOLECULE)	2ML
55397	IGG GOAT ANTI-HAMSTER IGG (WHOLE MOLECULE)	5ML
55400	FITC GOAT ANTI-HAMSTER IGG (WHOLE MOLECULE)	2ML
55402	HRP GOAT ANTI-HAMSTER IGG (WHOLE MOLECULE)	2ML
55933	PURIFIED HAMSTER IGG	20MG

  
Approved by: John Huang, PhD  
Quality Control Manager

**TO PLACE AN ORDER IN THE U.S., PLEASE CALL 800-854-0530  
OR TO OBTAIN TECHNICAL ASSISTANCE IN THE U.S., PLEASE CALL 800-279-5490  
OUTSIDE THE U.S., CONTACT YOUR LOCAL DISTRIBUTOR.**